

- 4 Johnson A, Wadsworth J, Wellings K, *et al.* *Sexual attitudes and lifestyles*. Oxford: Blackwell, 1994.
- 5 Adrien A, Leane V, Dassa C, *et al.* Sexual behaviour, condom use and HIV risk situations in the general population of Quebec. *Int J STD AIDS* 2001;12:108–15.
- 6 Laumann E, Gagnon J, Michael R, *et al.* *The social organisation of sexuality: sexual practices in the United States*. Chicago: University of Chicago Press, 1994.
- 7 Leridon H, van Zessen G, Hubert M. The Europeans and their sexual partners. In: Hubert M, Bajos N, Sandfort T, eds. *Sexual behaviour and HIV/AIDS in Europe*. London: UCL Press, 1998:165–96.
- 8 Konde-Lule JK, Sewankambo N, Morris M. Adolescent sexual networking and HIV transmission in rural Uganda. *Health Transit Rev* 1997;7(Suppl):89–100.
- 9 Morris M, Kretzschmar M. Concurrent partnerships and the spread of HIV. *AIDS* 1997;11:641–8.
- 10 Potterat JJ, Rothenberg RB, Muth SQ. Network structural dynamics and infectious disease propagation. *Int J STD AIDS* 1999;10:182–5.
- 11 Jolly AM, Muth SQ, Wyllie JL, *et al.* Sexual networks and sexually transmitted infections: a tale of two cities. *J Urban Health* 2001;78:433–45.
- 12 Smith AMA, Grierson J, Wain D, *et al.* Social network influences on gay men's sexual behaviour. *Sex Transm Infect* 2004;80:455–8.
- 13 Ellen JM, Brown BA, Chung SE, *et al.* Impact of sexual networks on risk for gonorrhea and chlamydia among low-income urban African American adolescents. *J Pediatr* 2005;146:518–22.
- 14 Rothenberg RB. The geography of gonorrhea. Empirical demonstration of core group transmission. *Sex Transm Infect* 1983;117:688–94.
- 15 Han Y, Coles FB, Muse A, *et al.* Assessment of a geographically targeted field intervention on gonorrhea incidence in two New York State counties. *Sex Transm Dis* 1999;26:296–302.
- 16 Zenilman JM, Ellish N, Fresia A, *et al.* The geography of sexual partnerships in Baltimore: applications of core theory dynamics using a geographic information system. *Sex Transm Dis* 1999;26:75–81.
- 17 Kandel DB, Kiros GE, Schaffran C, *et al.* Racial/ethnic differences in cigarette smoking initiation and progression to daily smoking: a multilevel analysis. *Am J Public Health* 2004;94:128–35.
- 18 Sundquist J, Malmström M, Johansson SE. Cardiovascular risk factors and the neighbourhood environment: a multilevel analysis. *Int J Epidemiol* 1999;28:841–5.
- 19 Subramanian SV, Chen JT, Rehkopf DH, *et al.* Racial disparities in context: a multilevel analysis of neighborhood variations in poverty and excess mortality among black populations in Massachusetts. *Am J Public Health* 2005;95:260–5.
- 20 Jun HJ, Subramanian SV, Gortmaker S, *et al.* A multilevel analysis of women's status and self-rated health in the United States. *J Am Med Womens Assoc* 2004;59:172–80.
- 21 Aral SO, Padian NS, Holmes KK. Advances in multilevel approaches to understanding the epidemiology and prevention of sexually transmitted infections and HIV: an overview. *J Infect Dis* 2005;191(Suppl 1):S1–6.
- 22 Diez Roux AV, Aiello AE. Multilevel analysis of infectious diseases. *J Infect Dis* 2005;191(Suppl 1):S25–33.
- 23 Leaver CA, Allman D, Meyers T, *et al.* Effectiveness of HIV prevention in Ontario, Canada: a multilevel comparison of bisexual men. *Am J Public Health* 2004;94:1181–5.
- 24 Browning CR, Leventhal T, Brooks-Gunn J. Neighborhood context and racial differences in early adolescent sexual activity. *Demography* 2004;41:697–720.
- 25 Cubbin C, Santelli J, Brindis CD, *et al.* Neighborhood context and sexual behaviors among adolescents: findings from the national longitudinal study of adolescent health. *Perspect Sex Reprod Health* 2005;37:125–34.
- 26 Smith AMA, Rissel CE, Richters J, *et al.* Sex in Australia: the rationale and methods of the Australian Study of Health and Relationships. *Aust N Z J Public Health* 2003;27:106–117.
- 27 Australian Bureau of Statistics. *Australian Standard Geographical Classification (ASGC) Cat no 1216.0*. Canberra: Australian Bureau of Statistics, 2001.
- 28 Australian Bureau of Statistics. *National Localities Index*, Cat no 1252.0. Canberra: Australian Bureau of Statistics, 2002.
- 29 Rasbash J, Steele F, Browne W, *et al.* *A user's guide to MLwiN*. London: University of London, 2004.
- 30 Glover J, Rosman D, Tennant S. Unpacking analyses relying on area-based data: are the assumptions supportable? *Int J Health Geogr* 2004;3:30.

COMMENTARY

Many sexual and reproductive health studies rely on individual level measures, especially those concerned with behavioural outcomes. However, it has been demonstrated in other fields of health research that area level factors matter, above and beyond individual level measures (see for example studies 17–20 cited by Smith and Subramanian¹). The need to simultaneously consider factors at individual and partnership

levels, as well as sexual networks and variations across subpopulations to better understand STI/HIV epidemiology has recently been emphasised.² The study by Smith and Subramanian thus provides important evidence in this respect.¹ The authors focus on numbers of heterosexual partners in the past year as their outcome but, as has been shown, partner numbers are an important predictor of a number of adverse sexual and reproductive health outcomes.^{3–4} Using data from the Australian Health and Relationship Study, a large national probability survey that achieved a high response rate and low item non-response, Smith and Subramanian report that area level factors, measured in terms of Australia's Statistical Sub Divisions, are significantly associated with partner numbers, even after taking account of individual level factors.

While the authors did not consider a particularly extensive range of area level factors, their data driven model may reflect difficulties in obtaining relevant aggregate level data consistent with the study's individual level measures and its spatial and temporal units of analysis. Of course, this assumes that concepts can be quantified and measured at the aggregate level, which is not always the case.^{5–6} These issues are not unique to this study and may hamper researchers' attempts to adopt a multilevel perspective. Despite these data challenges, multilevel analyses enable us to consider how factors operating at different levels are associated with each other, and in some cases, interact across the different levels of analysis. Indeed, it has been argued that this approach is not only more informative, but that just considering individual level "effects" may be erroneous as they may vary according to area level factors. Similarly, looking just at the aggregate level prevents distinction between individual "effects" and area "effects."⁷ (The use of quotation marks here is deliberate, reflecting how causality can not and should not be assumed from the analyses of cross sectional data, as Smith and Subramanian note.¹)

We should be encouraged to employ a more sophisticated, multilevel approach for examining sexual and reproductive health questions. The novel work by Smith and Subramanian inspires us to think beyond individual "effects" to advance our understanding of factors associated with sexual and reproductive risk behaviours and health outcomes. A more comprehensive understanding of both individual and area "effects" and their interaction will benefit geographically targeted interventions and increase the translational nature of sexual health research.

C H Mercer

Centre for Sexual Health & HIV Research, University College London,
Mortimer Market Centre, Off Capper Street, London WC1E 6JB, UK;
cmercer@gum.ucl.ac.uk

REFERENCES

- 1 Smith AMA, Subramanian SV. Population contextual associations with heterosexual partner numbers: a multilevel analysis. *Sex Transm Infect* 2006;82:250–4.
- 2 Aral SO, Padian NS, Holmes KK. Advances in multilevel approaches to understanding the epidemiology and prevention of sexually transmitted infections and HIV: an overview. *J Infect Dis* 2005;191(Suppl 1):S1–6.
- 3 Fenton KA, Korovessis C, Johnson AM, *et al.* Sexual behaviour in Britain: reported sexually transmitted infections and prevalent genital Chlamydia Trachomatis infection. *Lancet* 2001;358:1851–4.
- 4 Fenton KA, Mercer CH, Byron C, *et al.* Reported sexually transmitted disease clinic attendance and sexually transmitted infections in Britain: prevalence, risk factors, and proportionate population burden. *J Infect Dis* 2005;191(Suppl 1):S127–38.
- 5 Diez Roux AV, Aiello AE. Multilevel analysis of infectious diseases. *J Infect Dis* 2005;191(Suppl 1):S25–33.
- 6 Macintyre S, Ellaway A, Cummins S. Place effects on health: how can we conceptualise, operationalise and measure them? *Soc Soc Med* 2002;55:125–39.